

- Nagy, B. 1997. Orthoptera species and assemblages in the main habitat types of some urban areas in the Carpathian Basin. *Biologia* **52**: 233–240.
- Oly, M. 1996. Contribution to the Odonata in the urban area of Bochum – in 1993 and 1994. *Libellula* **15**: 11–26.
- Richter, K., Klausnitzer, B. & Zimdars, A. 1986. On the ant fauna of different urban-influenced ruderal places in the district of Leipzig (Hym., Formicidae). *Entomologische Nachrichten und Berichte* **30**: 115–120.
- Samways, M. J. & Steyler, N. S. 1996. Dragonfly (Odonata) distribution patterns in urban and forest landscapes and recommendations for riparian management. *Biological Conservation* **78**: 279–288.
- Sanderson, R. A. 1992. Hemiptera of naturally vegetated derelict land in north-west England. *Entomologist's Gazette* **43**: 221–226.
- Savage, A. A. 1989. Adults of the British aquatic Hemiptera Heteroptera: a key with ecological notes. *Freshwater Biological Association, Scientific Publication* **50**: 1–173.
- Schmitz, G. 1996. Urban ruderal sites as secondary habitats for phytophagous insects. *Verhandlungen der Gesellschaft für Ökologie* **26**: 581–585.
- Spalding, A. & Haes, E. C. M. 1995. Contaminated land—a resource for wildlife: a review and survey of insects on metalliferous mine sites in Cornwall. *Land Contamination and Reclamation* **3**: 24–29.
- Stewart, A. J. A. & Wright, A. F. 1995. A new inexpensive suction apparatus for sampling arthropods in grassland. *Ecological Entomology* **20**: 98–102.
- Wahlbrink, D. & Zucchi, H. 1994. Occurrence and settlement of carabid beetles on an urban railway embankment—a contribution to urban ecology. *Zoologische Jahrbücher, Abteilung für Systematik Ökologie und Geographie der Tier* **121**: 193–201.
- Wichtowska, M. & Sobczak, A. 1994. Formation of the water beetle (Coleoptera) fauna in conditions of the urban agglomeration of Szczecin (Western Pomerania). *Acta Hydrobiologica* **36**: 57–74.
- Williams, G. M. 1993. The colonisation of mining subsidence ponds by water boatman (Hemiptera: Heteroptera). *Entomologist's Gazette* **44**: 67–78.
- Wright, A. 1998. Hoverflies in a city environment: experiences in Coventry. *Dipterists Digest* **1**: 37–40.

SHORT COMMUNICATION

***Lasius brunneus* (Latr.) (Hymenoptera: Formicidae) found indoors.** – In May 2003, a naturalist colleague Mr Mike Paice gave me four ants; he had found them in a house in the village of Jacob's Well, near Guildford, Surrey, where they had been infesting a tin of biscuits. To my surprise, they were *Lasius brunneus*. In virtually all of the literature, this species is described as being 'fugitive', alluding to its evasive, secretive and non-aggressive habits. To find it invading a house is very unusual. From my own experience the normal ant found indoors is the ubiquitous black pavement ant, *Lasius niger* (L.), which commonly invades kitchens. Busy ant trails often lead to small amounts of spilled food in larders and eupboards or spoiled fruit at the bottom of the fruit bowl. In Norway and Sweden, *L. brunneus* is recorded from timbers in old houses and farm buildings (Collingwood, C. A. 1979. The Formicidae (Hymenoptera) of Fennoscandia and Denmark. *Fauna Ent. Scand.* **8**: 98–99). The house in Jacob's Well is old, part of it, and the adjoining barn having been built around 1640, so the ant may well have been nesting in some of the ancient building timber. – RICHARD A. JONES, 135 Friern Road, East Dulwich, London SE22 0AZ. bugmanjones@hotmail.com